

Antibiotic Allergy Cross Reactivity/Sensitivity Chart

Med you want to give →																					Allergic to ↓																		
Amikacin																				*																			
Amoxicillin																																							
Ampicillin	*																																						
Azithromycin																																							
Aztreonam																																							
Cefazolin																																							
Cefepime																																							
Cefotaxime																																							
Cefoxitin																																							
Ceftaroline																																							
Ceftazidime																																							
Ceftriaxone																																							
Cefuroxime																																							
Cephalexin	*	*																																					
Ciprofloxacin																																							
Clindamycin																																							
Daptomycin																																							
Doxycycline																																							
Ertapenem																																							
Erythromycin		*																																					
Gentamicin	*																																						
Imipenem																				*																			
Levofloxacin																		*																					
Linezolid																			*																				
Meropenem																			*																				
Metronidazole																				*																			
Moxifloxacin																			*																				
Penicillin																				*																			
Pip/tazo																					*																		
Sulfa																						*																	
Tetracycline																					*																		
Tigecycline																						*																	
Tobramycin	*																																						
Vancomycin																																							

Key: * = do NOT give | Blank = ok to give

This clinical tool is intended to be flexible. It serves as a reference point and/or recommendation, not rigid criteria. This clinical tool should be followed in most cases, but there is an understanding that, depending on the patient, setting, circumstances or factors, it can and should be tailored to fit individual needs.

Updated January 2019.